# Reliable quality assurance in beer production

## Detection of beer spoilage bacteria

**Products** 

#### **Key facts**

#### Simple

- Easy workflow with minimized hands-on-time
- No trained laboratory personnel needed

#### **Fast**

- Get your results faster
- No need to wait for external analysis results
- Always be one step ahead and make the right decisions on time

#### On-site

- Save time by doing the analysis at your site
- Always keep the full control from sampling to result
- No time-consuming shipping process of your samples



Early detection of beer spoilage is not only a matter of product safety, but also promotes sustainability. Endress+Hauser BioSense has developed a system to support customers in the rapid and early detection of beer spoilage bacteria and hop resistance genes. The system aims to assure product safety, compliance and sustainability.

#### Quality assurance is of paramount importance in beer production.

Beer spoilage bacteria have the potential to compromise the flavor, aroma, and overall quality of the beer. Implementing stringent quality control measures and monitoring systems ensures early detection and mitigation of such contaminants.

### The rapid detection of beer spoilers helps with

- Safeguarding the established flavor
- Maintaining customer trust
- Establishing a more sustainable production
- Minimisation of unusable batches and waste of material
- Using time & resources efficiently
- Optimisation of workflow
- Recognising problem areas in production
- Taking timely countermeasures

#### **Available Products**



Spoilage Bacteria Ident Identifies specific bacteria & hop resistance genes Parameters: L. brevis, horA & horC, expansion to 18 planned



Spoilage Bacteria Screen
Detects bacteria groups
& hop resistance genes
Parameters: Lactobacillus
& Pediococcus, Pectinatus,
Megasphaera, horA & horC



### Conventional PCR and Endress+Hauser BioSense workflow comparison

#### Step Conventional PCR workflow Endress+Hauser BioSense workflow Sampling and Manual or automated sampling at a customer-defined point in time enrichment Optional steps dependent on sample matrix and customer need: Homogenization Enrichment Closed system automating all steps Lysis ■ Two thermal devices necessary after sampling & enrichment High risk of mix-up and All reagents are pre-stored contamination while handling No manual intervention necessary buffers, enzymes and other reagents ■ 30 minutes hands-on time ■ 30 minutes lysis and inactivation DNA/RNA Centrifuge required Use of potentially hazardous purification chemicals High risk of mix-up and contamination while handling different reagents • 60 minutes hands-on time PCR and data Seperate lab rooms necessary Pipetting error-prone small volumes analysis 30 minutes hands-on time 60 minutes PCR duration Manual data interpretation required



Germany Sales V1.0

Endress+Hauser BioSense GmbH Georges-Köhler-Allee 302 79110 Freiburg im Breisgau Consultation Information Order

ehbs.endress.com

+491605135990 Info.ehbs@endress.com

